

2. Coselli JS, Lemaire SA, Koksoy C, Schmitling ZC, Curling PE. Cerebrospinal fluid drainage reduces paraplegia after thoracoabdominal aortic aneurysm repair: results of a randomized clinical trial. *J Vasc Surg.* 2002;35:631-9.
3. Estrera AL, Rubenstein FS, Miller CC 3rd, Huynh TT, Letsou GV, Safi HJ. Descending thoracic aortic aneurysm: surgical approach and treatment using the adjuncts cerebrospinal fluid drainage and distal aortic perfusion. *Ann Thorac Surg.* 2001;72:481-6.
4. Nienaber CA, Fattori R, Lund G, Dieckmann C, Walter W, von Kodolitsch Y, et al. Nonsurgical reconstruction of thoracic aortic dissection by stent graft placement. *N Engl J Med.* 1999;340:1539-45.
5. Mitchell RS, Miller DC, Dake MD, Semba CP, Moore KA, Sakai T. Thoracic aortic aneurysm repair with an endovascular stent-graft: the "first generation." *Ann Thorac Surg.* 1999;67:1971-4.
6. Crawford ES, Mizrahi EM, Hess KR, Coselli JS, Safi HJ, Patel VM. The impact of distal aortic perfusion and somatosensory evoked potential monitoring on prevention of paraplegia after aortic aneurysm operation. *J Thorac Cardiovasc Surg.* 1988;95:357-67.
7. Tiesenhausen K, Amann W, Koch G, Haugsegger KA, Oberwalder P, Rigler B. Cerebrospinal fluid drainage to reverse paraplegia after endovascular thoracic aortic aneurysm repair. *J Endovasc Ther.* 2000;7:132-5.

12/8/127787

doi:10.1067/mtc.2002.127787

### Reply to the Editor:

We appreciate the comments regarding our article by Oberwalder and colleagues and thank them for the valuable adjuncts regarding cerebrospinal fluid (CSF) drainage in stent-graft implantation.

Because of the restricted number of references in brief communications, we considered only publications in the high-impact journals. Nevertheless, we are aware of the report of Tiesenhausen and associates,<sup>1</sup> who described for the first time the impact of CSF drainage in a case of paraplegia after stent-graft implantation.

To close the entry tear in the descending aorta it is sometimes necessary to deploy a stent graft in the region (Th 8- L2) most prone to ischemia, but we agree that this should be avoided whenever possible. However, the risk of paraplegia has to be balanced against the risk of rupture of the aneurysm in these patients.

Routine use of CSF drainage in stent-graft implantations could help to reduce the risk of paraplegia in such cases.

Tatjana Fleck, MD

Martin Grabenwöger, MD

University of Vienna

Department of Cardiothoracic Surgery

Leitstelle 20A

Waehringer Guertel 18-20

Vienna, 1090 Austria

### Reference

1. Tiesenhausen K, Amann W, Koch G, Haugsegger KA, Oberwalder P, Rigler B. Cerebrospinal fluid drainage to reverse paraplegia after endovascular thoracic aortic aneurysm repair. *J Endovasc Ther.* 2000;7:132-5.

12/8/127793

### Alpha-stat strategy: Cause of ischemia in brains with old cerebral infarction despite selective cerebral hypothermic antegrade perfusion

#### To the Editor:

Washiyama and associates<sup>1</sup> are to be congratulated for experimentally verifying the anticipated susceptibility of brains with old infarction to become ischemic even during hypothermic selective continuous antegrade perfusion (SCP).

We do not doubt that SCP is better than no perfusion or retrograde perfusion, regardless of the pH management. However, despite SCP, ischemia developed in brains with old infarcts. Had the presented data been fully analyzed and interpreted without prejudice, the mechanisms of that ischemia would have been clear.

In our opinion, the alkalosis induced by alpha-stat hypothermic perfusion<sup>2</sup> explains every fact that the authors could not satisfactorily explain:

1. Although the authors failed to recognize and statistically analyze the increasing brain lactate efflux even in the control group from sample 1 to sample 4 during SCP, such increase seems to us significant. In our opinion, alpha-stat-induced alkalosis inhibits the creatine kinase catalyzed phosphorylation reactions.<sup>3</sup> This causes failure to aerobically synthesize high-energy~P bonds and switch to anaerobic metabolism with

consequent adenosine triphosphate consumption and lactate production even in the control group without prior infarct, despite the continuous perfusion.

2. Further lactate efflux increase in the infarct group after rewarming. Failure to use glucose aerobically becomes overt after rewarming following alpha-stat hypothermic uninterrupted perfusion at 20°C for 60 minutes.<sup>4</sup>

As pointed out by the authors, the penumbra area of an infarct is dependent on collateral flow. The vasoconstriction and decreased brain flow caused by alpha-stat strategies will certainly reduce such collateral flow, maximizing the metabolic effects of pH management. Sakamoto and associates<sup>5</sup> demonstrated brain anoxia development during early alpha-stat cooling before arrest. These effects are not limited to the period of cold perfusion but continue also during alpha-stat rewarming; thus, the lactate efflux at 32°C is maximal, even if circulatory arrest was not induced.

Obtaining the glutamate samples from the maxillary vein as often as the lactate samples would have answered whether the switch of aerobic to anaerobic metabolism and lactate efflux of the control group was enough to cause hypoxic excitotoxicity.<sup>6</sup>

pH-stat strategies increase brain blood flow and clearly should have been advantageous for perfusion of brains with old infarcts<sup>7</sup>; pH-stat hypothermic selective antegrade perfusion might prevent or minimize the metabolic effects of alpha-stat strategies even in brains with collateral flow-dependent penumbra areas.

Tadaomi-Alfonso Miyamoto, MD<sup>a</sup>

Koho-Julio Miyamoto, MD, PhD<sup>b</sup>

Research Department

Kokura Memorial Hospital

Kitakyushu-City, Japan<sup>a</sup>

Assistant Professor II Department of

Physiology

The University of the Ryukyus School of

Medicine

Okinawa, Japan<sup>b</sup>

### References

1. Washiyama N, Kazui T, Takinami M, Yamashita K, Fujita S, Terada H, et al. Experimental study on the effect of antegrade cerebral perfusion on brains with old cerebral infarction. *J Thorac Cardiovasc Surg.* 2001;122:734-40.